



THE VRAYLAR® SAVINGS PROGRAM

Eligible, commercially insured patients prescribed VRAYLAR may pay as little as \$0 for their 30-day or 90-day fills*

*Terms and conditions apply. See page 4 for program eligibility.

INDICATIONS AND USAGE

VRAYLAR (cariprazine) is indicated in adults as adjunctive therapy to antidepressants for the treatment of major depressive disorder (MDD), for the treatment of depressive episodes associated with bipolar I disorder (bipolar depression), for the acute treatment of manic or mixed episodes associated with bipolar I disorder, and for the treatment of schizophrenia.

Important Safety Information

WARNINGS: INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS and SUICIDAL THOUGHTS AND BEHAVIORS

- **Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. VRAYLAR is not approved for treatment of patients with dementia-related psychosis.**
- **Antidepressants increased the risk of suicidal thoughts and behaviors in pediatric and young adult patients in short-term studies. Closely monitor antidepressant-treated patients for clinical worsening, and for emergence of suicidal thoughts and behaviors. Safety and effectiveness of VRAYLAR have not been established in pediatric patients.**

Contraindication: VRAYLAR is contraindicated in patients with known hypersensitivity. Reactions have included rash, pruritus, urticaria, and reactions suggestive of angioedema.

Cerebrovascular Adverse Reactions, Including Stroke: In clinical trials with antipsychotic drugs, elderly patients with dementia had a higher incidence of cerebrovascular adverse reactions, including fatalities, vs placebo. VRAYLAR is not approved for the treatment of patients with dementia-related psychosis.

Neuroleptic Malignant Syndrome (NMS): NMS, a potentially fatal symptom complex, has been reported with antipsychotic drugs. NMS may cause hyperpyrexia, muscle rigidity, delirium, and autonomic instability. Additional signs may include elevated creatine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure. Manage with immediate discontinuation, intensive symptomatic treatment, and monitoring.

Please see additional Important Safety Information on the following pages. Please also see full [Prescribing Information](#), including Boxed Warnings, or visit https://www.rxabbvie.com/pdf/vraylar_pi.pdf

VRAYLAR Savings Program— Financial Support for Your Patients



With the VRAYLAR Savings Card your prescribed patients may pay:

\$0 FOR 30-DAY OR 90-DAY PRESCRIPTION

Eligible, commercially insured patients prescribed VRAYLAR may pay as little as **\$0 for 30-day or 90-day fills** of VRAYLAR.*

\$0 FOR A GENERIC ANTIDEPRESSANT

Eligible, commercially insured patients prescribed VRAYLAR as an adjunctive treatment for major depressive disorder may pay as little as **\$0** for their generic antidepressant.*

3 WAYS PATIENTS CAN ENROLL

1. Text VRAYLAR to 20866[†]



2. Visit vraypay.com



3. Scan QR code



*Terms and conditions apply. See page 4 for program eligibility.

[†]Message and data rates may apply. See page 4 for details.

Important Safety Information (continued)

Tardive Dyskinesia (TD): Risk of developing TD (a syndrome of potentially irreversible, involuntary, dyskinetic movements) and the likelihood it will become irreversible may increase with the duration of treatment and the cumulative dose. The syndrome can develop after a relatively brief treatment period, even at low doses, or after treatment discontinuation. If signs and symptoms of TD appear, drug discontinuation should be considered.


Late-Occurring Adverse Reactions: Adverse reactions may first appear several weeks after initiation of VRAYLAR, probably because plasma levels of cariprazine and its major metabolites accumulate over time. As a result, the incidence of adverse reactions in short-term trials may not reflect the rates after longer-term exposures. Monitor for adverse reactions, including extrapyramidal symptoms (EPS) or akathisia, and patient response for several weeks after starting VRAYLAR and after each dosage increase. Consider reducing the dose or discontinuing the drug.

Please see additional Important Safety Information on the following pages. Please also see full [Prescribing Information](#), including Boxed Warnings, or visit https://www.rxabbvie.com/pdf/vraylar_pi.pdf

Patients With Coverage Restrictions May Be Eligible for the VRAYLAR Savings Program


For patients experiencing access challenges:

The VRAYLAR Savings Program may help eligible patients **pay as little as \$40** for their VRAYLAR prescriptions.*



Up to **two**
30-day fills

if payers have yet to determine coverage for VRAYLAR



Up to **two**
30-day fills

if a prior authorization or step therapy is required



Have questions about the VRAYLAR Savings Program?
Contact your **Field Reimbursement Manager for support.**

*Terms and conditions apply. See page 4 for program eligibility.

Important Safety Information (continued)

Metabolic Changes: Atypical antipsychotics, including VRAYLAR, have caused metabolic changes, such as:

- **Hyperglycemia and Diabetes Mellitus:** Hyperglycemia, in some cases associated with ketoacidosis, hyperosmolar coma, or death, has been reported in patients treated with atypical antipsychotics. Assess fasting glucose before or soon after initiation of treatment, and monitor periodically during long-term treatment.
- **Dyslipidemia:** Atypical antipsychotics cause adverse alterations in lipids. Before or soon after starting an antipsychotic, obtain baseline fasting lipid profile and monitor periodically during treatment.
- **Weight Gain:** Weight gain has been observed with VRAYLAR. Monitor weight at baseline and frequently thereafter.

Please see additional Important Safety Information on the following page. Please also see full [Prescribing Information](#), including Boxed Warnings, or visit https://www.rxabbvie.com/pdf/vraylar_pi.pdf

Important Safety Information (continued)

Leukopenia, Neutropenia, and Agranulocytosis: Leukopenia/neutropenia have been reported with antipsychotics, including VRAYLAR. Agranulocytosis (including fatal cases) has been reported with other antipsychotics. Monitor complete blood count in patients with pre-existing low white blood cell count (WBC)/absolute neutrophil count or history of drug-induced leukopenia/neutropenia. Discontinue VRAYLAR at the first sign of a clinically significant decline in WBC and in severely neutropenic patients.

Orthostatic Hypotension and Syncope: Atypical antipsychotics cause orthostatic hypotension and syncope, with the greatest risk during initial titration and with dose increases. Monitor orthostatic vital signs in patients predisposed to hypotension and in those with cardiovascular/cerebrovascular diseases.

Falls: VRAYLAR may cause somnolence, postural hypotension, motor and sensory instability, which may lead to falls and, consequently, fractures or other injuries. For patients with diseases, conditions, or medications that could exacerbate these effects, complete fall risk assessments when initiating antipsychotics and recurrently for patients on long-term therapy.

Seizures: Use VRAYLAR with caution in patients with a history of seizures or with conditions that lower the seizure threshold.

Potential for Cognitive and Motor Impairment: Somnolence was reported with VRAYLAR. Caution patients about performing activities requiring mental alertness (eg, operating hazardous machinery or a motor vehicle).

Body Temperature Dysregulation: Use VRAYLAR with caution in patients who may experience conditions that increase body temperature (eg, strenuous exercise, extreme heat, dehydration, or concomitant anticholinergics).

Dysphagia: Esophageal dysmotility and aspiration have been associated with antipsychotics. Antipsychotic drugs, including VRAYLAR, should be used cautiously in patients at risk for aspiration.

Drug Interactions: Strong and moderate CYP3A4 inhibitors increase VRAYLAR concentrations, so VRAYLAR dose reduction is recommended. Concomitant use with CYP3A4 inducers is not recommended.

Adverse Reactions: The most common adverse reactions in clinical trials ($\geq 5\%$ and at least twice the rate of placebo) are listed below:

- **Adjunctive Treatment of Major Depressive Disorder:**

- In 6-week, fixed-dose trials the incidences within the recommended doses (VRAYLAR 1.5 mg/day + antidepressant therapy [ADT] or 3 mg/day + ADT vs placebo + ADT) were akathisia (7%, 10% vs 2%), nausea (7%, 6% vs 3%), and insomnia (9%, 10% vs 5%).

- In one 8-week flexible-dose trial, incidences within the doses (VRAYLAR 1-2 mg/day + antidepressant therapy [ADT] or 2-4.5 mg/day + ADT vs placebo + ADT) were akathisia (8%, 23% vs 3%), restlessness (8%, 8% vs 3%), fatigue (7%, 10% vs 4%), constipation (2%, 5% vs 2%), nausea (7%, 13% vs 5%), increased appetite (2%, 5% vs 2%), dizziness (4%, 5% vs 2%), insomnia (14%, 16% vs 8%), and extrapyramidal symptoms (12%, 18% vs 5%).

- **Bipolar Mania:** The incidences within the recommended dose range (VRAYLAR 3–6 mg/day vs placebo) were EPS (26% vs 12%), akathisia (20% vs 5%), vomiting (10% vs 4%), dyspepsia (7% vs 4%), somnolence (7% vs 4%), and restlessness (7% vs 2%).

- **Bipolar Depression:** The incidences within the recommended doses (VRAYLAR 1.5 mg/day or 3 mg/day vs placebo) were nausea (7%, 7% vs 3%), akathisia (6%, 10% vs 2%), restlessness (2%, 7% vs 3%), and EPS (4%, 6% vs 2%).

- **Schizophrenia:** The incidences within the recommended dose range (VRAYLAR 1.5–3 mg/day and 4.5–6 mg/day vs placebo) were EPS (15%, 19% vs 8%) and akathisia (9%, 13% vs 4%).

Please also see full **Prescribing Information**, including **Boxed Warnings**, or visit

https://www.rxabbvie.com/pdf/vraylar_pi.pdf

*Eligibility: Available to patients with commercial insurance coverage for VRAYLAR[®] who meet eligibility criteria. This co-pay assistance program is not available to patients receiving prescription reimbursement under any federal, state, or government-funded insurance programs (for example, Medicare [including Part D], Medicare Advantage, Medigap, Medicaid, TRICARE, Department of Defense, or Veterans Affairs programs) or where prohibited by law. Offer subject to change or termination without notice. Restrictions, including monthly maximums, may apply. This is not health insurance. **For full Terms and Conditions**, visit [VRAYLARSavingscard.com](https://www.vraylarsavingscard.com) or call 1-800-761-0436 for additional information. To learn about AbbVie's privacy practices and your privacy choices, visit <https://abbv.ie/corpprivacy>

*Message and data rates may apply. You are not required to consent as a condition of receiving goods or services. You can reply HELP for help. You can reply STOP to opt out at any time. By texting VRAYLAR to 20686, you agree to AbbVie's Mobile Terms and Conditions at <https://abbv.ie/USMobileTerms> and Privacy Notice at <https://abbv.ie/corpprivacy>. Full Prescribing Information available at https://www.rxabbvie.com/pdf/vraylar_pi.pdf.